

What You Need To Know About Your Medicine For Tuberculosis (TB):

Isoniazid (INH) Oral

About Your Medicine:

ISONIAZID (eye-soe-NYE-a-zid) is an antibiotic used to prevent or treat tuberculosis (TB).

Before Using This Medicine:

Tell your doctor, nurse, and pharmacist if you:

- Have drug allergies
- Are pregnant or plan to become pregnant
- Are breastfeeding
- Are taking any other prescription medicines
- Are taking over-the-counter medicines (such as cold medicines, pain relievers, vitamins, herbs, etc.)
- Have any other medical problems including diabetes, gout, liver disease, or kidney disease

Tips to Help You Take Your Medicine:

- Take exactly as directed.
- Do not miss any doses or take two doses at the same time.
- Do not share your medicines with others.
- Keep this and all other medicines out of the reach of children.
- Store in a cool, dry, place away from extreme heat or cold.

www.scdhec.gov/tb

Warnings:

- DO NOT drink alcohol while on this medicine to avoid possible liver problems.
- If you think you have taken an overdose of this medicine, call Poison Control at 1-800-222-1222.

Possible Side Effects of TB Medicines:

- Upset stomach or stomach pain
- Nausea or vomiting
- Severe weakness or tiredness
- Flu-like symptoms
- Coffee/tea colored urine or light stools
- Yellow eyes or skin
- Rash or itching
- Easily bleeding or bruising
- Blurred vision
- Pain or tingling in hands, feet, or joints
- Dizziness
- Fever

INH may:

- Cause dizziness
- Cause a tingling sensation or numbness of mouth, hands, and feet



Please contact your DHEC nurse immediately if you experience any of these side effects.

Những Gì Quý Vị Cần Biết về Thuốc Điều Trị Bệnh Lao (TB) của Quý Vị:

Isoniazid (INH) Dùng Đường Uống

Giới Thiệu về Thuốc của Quý Vị:

ISONIAZID (eye-soe-NYE-a-zid) là một loại kháng sinh được sử dụng để ngăn ngừa hoặc điều trị bệnh lao (TB).

Trước Khi Sử Dụng Loại Thuốc Đây:

Hãy cho bác sĩ, y tá, và dược sĩ của quý vị biết nếu quý vị:

- Bị dị ứng thuốc
- Đang mang thai hoặc dự định có thai
- Đang cho con bú
- Đang dùng bất kỳ loại thuốc theo toa nào khác
- Đang dùng những loại thuốc không theo toa (như thuốc cảm cúm, thuốc giảm đau, vitamin, thảo dược, v.v.)
- Có bất kỳ vấn đề y tế nào khác bao gồm bệnh tiểu đường, bệnh gout, bệnh gan hoặc bệnh thận

Lời Khuyên Giúp Quý Vị Dùng Thuốc:

- Dùng đúng theo chỉ dẫn.
- Không bỏ lỡ bất kỳ liều nào hay dùng hai liều cùng một lúc.
- Không chia sẻ thuốc của quý vị với người khác.
- Giữ loại thuốc này và tất cả các loại thuốc khác ngoài tầm với của trẻ em.
- Bảo quản ở nơi mát mẻ, khô ráo, tránh xa nhiệt độ quá nóng hoặc quá lạnh.

www.scdhec.gov/tb

Cảnh báo:

- KHÔNG uống rượu bia trong khi dùng loại thuốc này để tránh các vấn đề gan có thể xảy ra.
- Nếu quý vị cho rằng mình đã dùng loại thuốc này quá liều, hãy gọi cho Bộ Phận Kiểm Soát Chất Độc theo số 1-800-222-1222.

Tác Dụng Phụ Có Thể Có của Thuốc Điều Trị Bệnh Lao (TB):

- Rối loạn dạ dày hoặc đau dạ dày
- Buồn nôn hoặc nôn mửa
- Rất yếu hoặc mệt mỏi
- Các triệu chứng giống như cúm
- Nước tiểu có màu cà phê/trà hoặc phân có màu nhạt
- Vàng mắt hoặc vàng da
- Phát ban hoặc ngứa
- Dễ chảy máu hoặc bầm tím
- Mờ mắt
- Đau hoặc ngứa ran ở bàn tay, bàn chân, hoặc khớp
- Chóng mặt
- Sốt

INH có thể:

- Gây chóng mặt
- Gây cảm giác ngứa ran hoặc tê miệng, bàn tay và bàn chân



Hãy liên lạc với y tá DHEC của quý vị ngay lập tức nếu quý vị gặp bất kỳ tác dụng phụ nào trong số này.